

LK91713

Biotin-Linked Semaphorin 3A (SEMA3A)



Order 021-34695924
orders@ab-mart.com
Support 400-6123-828
support1@ab-mart.com
Web www.ab-mart.com.cn


□ 100ul;

UniProt : O08665

Alternative Names:

SEMAL; Hsema-I; Hsema-III; SEMA1; SEMAD; SEMAIII; SemD; Coll-1; Sema Domain, Immunoglobulin Domain(Ig), Transmembrane Domain(TM) and Short Cytoplasmic Domain 3A

Application:



Reactivity : Mouse

Application : WB IHC IF

Dilution:

Western blotting: 0.5-2ug/ml

Immunocytochemistry in formalin fixed cells: 5-20ug/ml

Immunohistochemistry in formalin fixed frozen section: 5-20ug/ml

Immunohistochemistry in paraffin section: 5-20ug/ml

Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml

Optimal working dilutions must be determined by end user.

Purification:

Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Storage Condition and Buffer:

0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

Tag : CD & Adhesion molecule;

Form : Liquid

Concentration : 500µg/mL